

Karyotype Analysis of Five *Bellevallia* Lapeyr. Species (Asparagaceae) from Turkey

Hüseyin Eroğlu¹, Süleyman Mesut Pınar², Gulden Dogan^{3*},
M. Yavuz Paksoy⁴ and Yasar Kiran³

¹ Van Yüzüncü Yıl University, Faculty of Sciences, Department of Biology, Tuşba-Van, Turkey

² Van Yüzüncü Yıl University, Van School of Health, Tuşba-Van, Turkey

³ Firat University, Science Fac., Biology Department, Elazığ, Turkey

⁴ Department of Environmental Engineering, Faculty of Engineering, Munzur University, Tunceli, Turkey

Received May 24, 2022; accepted June 15, 2022

Summary The chromosome numbers and karyotypes of five species of the genus *Bellevallia* distributed in Turkey were analyzed. These taxa are *B. anatolica* B. Mathew & Özhatay, *B. fominii* Woronow, *B. kurdistanica* Feinbrun, *B. rixii* Wendelbo, *B. turcica* Pınar & Eroğlu. Three of them (*B. anatolica*, *B. rixii* and *B. turcica*) are endemic in Turkey. The chromosome numbers of all studied taxa were determined as $2n=2x=8$. Haploid chromosome lengths varied from $57.78\mu\text{m}$ (*B. fominii*) to $68.17\mu\text{m}$ (*B. anatolica*) among species. Karyotype analysis showed that *Bellevallia* taxa generally have median (m), submedian (sm), and subterminal (st) chromosomes, but not subterminal chromosomes in *B. fominii* species.

Keywords Asparagaceae, *Bellevallia*, Chromosome number, Karyotype, Turkey.

The genus *Bellevallia* is distributed on three continents: Asia, Europe, and Africa. The continent with the richest number of species in Asia. Especially Anatolia and Iran, as well as Transcaucasia, Palestine, and Syria are the areas where the *Bellevallia* genus is distributed. Considering the numerical distribution of the genus *Bellevallia*, 35 species are distributed in Asia, eight species in Africa, and seven species in Europe. In addition, 38 species are found in the Irano-Turanian, 16 species in the Mediterranean, and one species in the Saharo-Sindian Phytogeographic region (Feinbrun 1939). In the monograph study on the genus *Bellevallia*, it was stated that 14 of the 45 taxa were distributed in Anatolia (Feinbrun 1940). According to Wendelbo (1984) in Turkey, 18 species, seven of which are endemic; According to the List of Plants of Turkey, it is 24 (Tugay 2012); There are 35 species in total, with new species identified in recent years. (Uzunhisarcıklı *et al.* 2013, Gürdal *et al.* 2014, Yıldırım *et al.* 2014, Karabacak *et al.* 2014, 2015, Pınar *et al.* 2016, Şahin *et al.* 2016, Fidan 2019, Yıldırım and Sefalı 2020).

The number, size, and shape of chromosomes can be used to characterize the karyotype of plants (Jahan *et al.* 2014). The genus *Bellevallia* is characterized by a basic chromosome number ($x=4$). Most representatives

of the genus have a set of $2n=2x=8$ chromosomes in the complement. Chromosome complement in *Bellevallia* seems to be stable, having a typical karyotype occurring in almost all species studied. However, polyploidy has also been observed in some studies. According to the literature, it has also been determined that some species of this genus are triploid, tetraploid and hexaploid (Özhatay *et al.* 1991a, b, Johnson 2003, Persson 2006, Yaylacı *et al.* 2009, Uzunhisarcıklı *et al.* 2013, Gürdal *et al.* 2014, Karabacak *et al.* 2014, 2015, Pınar *et al.* 2016).

This paper aims to clearly define the chromosome numbers and morphology of the five *Bellevallia* species and to bring the karyotype data to the literature.

Materials and methods

Plant materials of five *Bellevallia* species were collected from natural habitats during the fruiting season between 2019 and 2021. The species name, localities, geographical position, altitude, and voucher number are presented in Table 1. The voucher specimens were de-

Table 1. Localities and voucher numbers of the studied *Bellevallia* taxa.

Taxa	Locality	Voucher number
<i>B. anatolica</i>	B7/Tunceli: Tüllük Village, 1435 m	Eroğlu, 1666
<i>B. fominii</i>	B9/Van: Gevaş, 1887 m	Eroğlu, 1680
<i>B. kurdistanica</i>	B7/Tunceli: Ovacık, 1400 m	Dogan, 2590
<i>B. rixii</i>	B9/Van: Başkale, Güzeldere, 2632 m	Eroğlu, 1610
<i>B. turcica</i>	C5/Adana: Pozanti, 1515 m	Pınar, 6502

* Corresponding author, e-mail: gdogan@firat.edu.tr

DOI: 10.1508/cytologia.87.319

posited at the Herbarium of Firat University.

The karyological studies are conducted on the meristematic cells of seven bulbs of each species. The bulbs were germinated at 25°C. The actively growing root tips were pretreated with 0.05% colchicine for 4 h at room temperature. Afterward, the root tips were fixed with Carnoy fixative (1:3 glacial acetic acid–absolute ethanol) for at least 24 h at 4°C, hydrolyzed in 1 M HCl at 60°C for 10–15 min, then rinsed in tap water for 3–5 min. Finally, they were stained in the Feulgen reagent for 1 h and mounted in 45% acetic acid (Kiran *et al.* 2012). Digital microphotographs from at least five well-spread metaphase plates were taken using an Olympus BX51 microscope and an Olympus Camedia C-4000 digital camera.

The number of somatic chromosomes, chromosome length range, haploid chromosome length, arm ratio, and

relative length were measured. The karyotype formula was determined based on the centromere position using a system of Levan *et al.* (1964). The ideograms of these taxa are arranged in decreasing lengths according to the chromosome size in the metaphase (Martin *et al.* 2009).

Results and discussion

The chromosome number, chromosome length range, haploid chromosome length, arm ratio, relative length, and karyotype formula were determined from 14 metaphase cells in each species and given in Table 2. Metaphase chromosomes are shown in Fig. 1 and ideograms in Fig. 2.

The somatic chromosome number of *B. anatolica* was $2n=2x=8$ (Fig. 1A). The karyotype formula

Table 2. Karyological features of the studied *Bellevia* taxa.

Taxa	$2n$	Chromosome length (μm)		Haploid chromosome length (μm)	Arm ratio		Relative length (%)		Karyotype formula
		Min	Max		Min	Max	Min	Max	
<i>B. anatolica</i>	8	13.79	22.41	68.17	1.24	4.00	20.22	32.88	$2m+sm+st$
<i>B. fominii</i>	8	10.19	17.43	57.78	1.12	2.76	19.31	33.02	$2m+2sm$
<i>B. kurdistanica</i>	8	11.67	20.00	59.10	1.33	3.91	19.75	33.83	$2m+sm+st$
<i>B. rixii</i>	8	12.05	18.82	60.68	1.23	3.40	19.86	31.02	$2m+sm+st$
<i>B. turcica</i>	8	11.96	20.17	61.88	1.49	3.78	19.32	32.60	$m+2sm+st$




Fig. 1. Metaphase chromosomes of studied *Bellevia* taxa. A. *B. anatolica*, B. *B. fominii*, C. *B. kurdistanica*, D. *B. rixii*, E. *B. turcica* Scale bars=10 μm .




Fig. 2. Haploid idiograms of studied *Bellevia* taxa. A. *B. anatolica*, B. *B. fominii*, C. *B. kurdistanica*, D. *B. rixii*, E. *B. turcica*.

was $2m+sm+st$. The chromosome length ranged from 13.79 to 22.41 μm and the haploid chromosome length was 68.17 μm . Arm ratios were 1.24–4.00 and relative lengths were 20.22–32.88% (Table 2). The idiogram is given in Fig. 2A. The somatic chromosome number of *B. fominii* was $2n=2x=8$ (Fig. 1B). The karyotype formula was $2m+12sm$. The chromosome length ranged from 10.19 to 17.43 μm and the haploid chromosome length was 57.78 μm . Arm ratios were 1.12–2.76 and relative lengths were 19.31–33.02% (Table 2). The idiogram is given in Fig. 2B. The somatic chromosome number of *B. kurdistanica* was $2n=2x=8$ (Fig. 1C). The karyotype formula was $2m+sm+st$. The chromosome length ranged from 11.67 to 20.00 μm and the haploid chromosome length was 59.10 μm . Arm ratios were 1.33–3.91 and relative lengths were 19.75–33.83% (Table 2). The idiogram is given in Fig. 2C. The somatic chromosome number of *B. rixii* was $2n=2x=8$ (Fig. 1D). The karyotype formula was $2m+sm+st$. The chromosome length ranged from 12.05 to 18.82 μm and the haploid chromosome length was 60.68 μm . Arm ratios were 1.23–3.40 and relative lengths were 19.86–31.02% (Table 2). The idiogram is given in Fig. 2D. The somatic chromosome number of *B. turcica* was $2n=2x=8$ (Fig. 1E). The karyotype formula was $m+2sm+st$. The chromosome length ranged from 11.96 to 20.17 μm and the haploid chromosome length was 61.88 μm . Arm ratios were 1.49–3.78 and relative lengths were 19.32–32.60% (Table 2). The idiogram is given in Fig. 2E.

Chromosome lengths show wide variation among species. The mean chromosome length varies between 10.19 μm (*B. fominii*) and 22.41 μm (*B. anatolica*). Haploid chromosome length ranges from 57.78 μm (*B. fominii*) to 68.17 μm (*B. anatolica*). The arm ratio was determined minimum of 1.12 in *B. fominii*, and maximum of 4.00 in *B. anatolica*, whereas the lowest relative length was observed in *B. fominii* (19.31%) and the highest relative length was found in *B. kurdistanica* (33.83%).

To date, only one study has been conducted on the chromosome number of the *B. anatolica*. In this study conducted by Johnson (1994), the chromosome number of the species was reported as $2n=8$. Our investigations also determined the same $2n=8$. Numerous studies have

been conducted on *B. fominii* and the somatic chromosome number of this species is generally reported as $2n=8$ (Delaunay 1926, Fedorov 1969, Özhatay *et al.* 1991a, b, Johnson and Brandham 1997, Bothmer and Wendelbo 1981, Nersesian 2001, Johnson 2003).

According to the literature, the chromosome number of *B. kurdistanica* species has been reported as $2n=8$, 16 (Al-Mudarris 1973, Bothmer and Wendelbo 1981, Johnson and Brandham 1997). The somatic chromosome number of *B. rixii* has been reported as $2n=8$ (Özhatay *et al.* 1991a, b, Johnson 2003). The chromosome number of *B. turcica*, which was introduced to the scientific world as a new species for Turkey by Pınar and Eroğlu (2016), was determined as $2n=8$ in the same publication.

Although the genus *Bellevia* is closely related to *Muscari*, *Hyacinthus*, and *Hyacinthella*, sharing with them considerable morphological similarity, its karyotype is unique and distinguishable. The 'basic karyotype' of *Bellevia* consists of $x=4$, with long chromosomes (Bothmer and Wendelbo 1981). With this study, we aim to contribute to the search for karyologically understudied species to expand the karyological knowledge of the genus *Bellevia*.

References

- Al-Mudarris, H. E. 1973. Cytological studies on the genus *Bellevia* Lapeyr from Iraq. Bull. Coll. Sci. Univ. Baghdad **14**: 177–186.
- Bothmer, R. V. and Wendelbo, P. 1981. Cytological and morphological variation in *Bellevia*. Nord. J. Bot. **1**: 4–11.
- Delaunay, L. 1926. La theorie chromosomique de l'heredite et les chromosomes chez quelques Liliacees. Vist. Tiflis. Bot. Sada N.S. **2**
- Fedorov, A. A. (ed.) 1969. Chromosome Numbers of Flowering Plants. Komarov V.L. Botanical Institute, Leningrad.
- Feinbrun, N. 1939. A monographic study on the genus *Bellevia* Lapeyr. Palestine J. Bot. **1**: 131–142.
- Feinbrun, N. 1940. A monographic study on the genus *Bellevia* Lapeyr. Palestine J. Bot. **1**: 336–409.
- Fidan, M. 2019. *Bellevia sasonii* (Asparagaceae): A new species from Turkey. Phytotaxa **394**: 126–132.
- Gürdal, B., Özhatay, N., Koçyiğit, M. and Kaya, E. 2014. Two New Species for Turkish Geophyte Flora in the Genus *Bellevia*. In: Kaya, E. (ed.) Turkey Geophytes, vol. 3. Atatürk Horticultural Central Research Institute, Yalova. pp. 525–531. (in Turkish).
- Jahan, B., Vahidy, A. A., Saeed, R. and Mirbahar, A. A. 2014. Karyological studies in ten different populations of desert lily *Aloe vera* from Pakistan. Pak. J. Bot. **46**: 1731–1734.
- Johnson, M. A. T. 1994. Cytology of three new geophytes from Turkey. Kew Bull. **49**: 491–498.
- Johnson, M. A. T. 2003. Polyploidy and karyotype variation in Turkish *Bellevia* (Hyacinthaceae). Bot. J. Linn. Soc. **143**: 87–98.
- Johnson, M. A. T. and Brandham, P. E. 1997. New chromosome numbers in petaloid monocotyledons and in other miscellaneous angiosperms. Kew Bull. **52**: 121–138.
- Karabacak, O., Yıldırım, H. and Martin, E. 2014. *Bellevia pseudolongipes* sp. nov. (Asparagaceae): A new species from south eastern Anatolia, Turkey. Phytotaxa **188**: 209–217.
- Karabacak, O., Yıldırım, H. and Martin, E. 2015. *Bellevia koyuncui* sp. nova (Asparagaceae): A new species from south eastern

- Anatolia, Turkey. *Phytotaxa* **203**: 81–84.
- Kiran, Y., Turkoglu, I., Kirilmaz, F., Arabaci, T., Sahin, A. and Bagci, E. 2012. Karyological investigation of six *Achillea* L. (Asteraceae) species growing in Turkey. *Caryologia* **65**: 101–105.
- Levan, A., Fredga, K. and Sandberg, A. 1964. Nomenclature of centromeric position of chromosomes. *Hereditas* **52**: 201–220.
- Martin, E., Dinç, M. and Duran, A. 2009. Karyomorphological study of eight *Centaurea* L. taxa (Asteraceae) from Turkey. *Turk. J. Bot.* **33**: 97–104.
- Nersesian, A. 2001. A karyosystematic study of Armenian *Muscari* and *Bellevia* (Hyacinthaceae). *Bocconea* **13**: 383–389.
- Özhatay, N., Johnson, M. A. T. and Mathew, B. 1991a. Chromosome numbers of Turkish *Bellevia* species, including a new hexaploid from European Turkey. *Bot. Chron.* **10**: 813–818.
- Özhatay, N., Johnson, M. A. T., Mathew, B. and Dalgıç, G. 1991b. A new hexaploid *Bellevia* (Hyacinthaceae) from European Turkey. *Bot. J. Linn. Soc.* **107**: 89–99.
- Persson, K. 2006. One new and one emended species of *Bellevia* (Hyacinthaceae) from Turkey. *Bot. J. Linn. Soc.* **150**: 253–260.
- Pınar, S. M., Eroğlu, H. and Fidan, M. 2016. *Bellevia behcetii* sp. nov. (Asparagaceae): A new species from south eastern Anatolia, Turkey. *Phytotaxa* **270**: 127–136.
- Şahin, B., Aslan, S., Karabacak, O. and Martin, E. 2016. *Bellevia vuralii* B. Şahin & Aslan (Asparagaceae): A new species from SE Turkey. *Turk. J. Bot.* **40**: 394–401.
- Tugay, O. 2012. *Bellevia* Lapeyr. In: Güner, A., Aslan, S., Ekim, T., Vural, M. and Babaç, M. T. (eds). List of Plants of Turkey, Nezahat Gökyiğit Botanical Garden and Flora Research Association Publication, Istanbul. pp. 95–96. (*in Turkish*).
- Uzunhisarcıklı, M. E., Duman, H. and Yılmaz, S. 2013. A new species of *Bellevia* (Hyacinthaceae) from Turkey. *Turk. J. Bot.* **37**: 651–655.
- Wendelbo, P. 1984. *Bellevia* Lapeyr. In: Davis, P. H., Mill, R. R. and Tan, K. (eds). Flora of Turkey and the East Aegean Islands, Vol. 8. Edinburgh University Press, Edinburgh. pp. 264–274.
- Yaylacı, Ö. K., Koyuncu, O., Öztürk, D. and Tokur, S. 2009. *Bellevia clusiana* Griseb. (Hyacinthaceae) cytotoxic characteristics. *Cankaya Univ. J. Sci. Engineb.* **12**: 193–200. (*in Turkish*)
- Yıldırım, H. and Sefalı, A. 2020. *Bellevia bayburtensis* (Asparagaceae): A new species from northeastern Anatolia, Turkey. *Phytotaxa* **441**: 285–293.
- Yıldırım, H., Altıoğlu, Y., Şahin, B. and Aslan, S. 2014. *Bellevia chrisii* sp. nov. (Asparagaceae) from eastern Anatolia, Turkey. *Nord. J. Bot.* **33**: 45–49.
-